



UCL



Developmental
Risk and Resilience Unit

Childhood maltreatment,
latent vulnerability and the
shift to preventative help:

*Understanding the link between
childhood maltreatment and
long-term mental health risk*

Eamon McCrory PhD DCLinPsy

Professor of Developmental Neuroscience &
Psychopathology, UCL

e.mccrory@ucl.ac.uk



Anna Freud
National Centre for
Children and Families

Coram Kent 23rd March 2018


Poor
outcomes

- ↑ Psychiatric disorders
- ↓ Attainment
- ↓ Economic productivity
- ↓ Physical Health

Adversity



Healthy development



Infancy.....Childhood.....Adolescence.....Adulthood

Poor
outcomes

↑ Psychiatric
disorders

Adversity





- May emerge many years later
- More likely to be less responsive to traditional treatments
- Problems more likely to be comorbid
- Problems show greater severity



Infancy.....Childhood.....Adolescence.....Adulthood

Maltreatment
Exposure



Treatment

Resilient
outcome

Limitations of a psychiatric diagnostic model:

- Focus on trauma can obscure other mental health needs (Green et al., 2016; Woolgar et al., 2015)
- Focus on diagnostic categories can obscure causal factors and relevant situational factors.
- Mental health is viewed in terms of cut-offs rather than as a continuum. Problems often reach crisis point before they are addressed.
- Often an absence of a child-centered needs-orientated approach that seeks to understand the child's presentation in a holistic way.
- As a result children often do not get the right help in a timely way as many CAMHS professionals feel ill-equipped to deal with the complexity that the child (and system around them) presents with.


Adversity



Poor
outcome

Resilient
outcome

Adversity



Poor
outcome

Resilient
outcome

1. **Early prevention**— how can we better help and support child who have experienced maltreatment to prevent the emergence of later problems?
2. **Pinpointing mechanisms** – how does adversity get under the skin and can an understanding of this help us think about preventative approaches?

The concept of Latent Vulnerability

McCrory & Viding

Development and Psychopathology, 2015

Development and Psychopathology 27 (2015), 493–505
© Cambridge University Press 2015
doi:10.1017/S0954579415000115

The theory of latent vulnerability: Reconceptualizing the link
between childhood maltreatment and psychiatric disorder

Adversity




?





Poor
outcome

Resilient
outcome


- Markers of latent vulnerability should be associated with maltreatment experience
- They are not necessarily symptoms
- They should be present even in the absence of psychiatric disorder
- They should be predictive of *future* psychiatric risk







Adversity



Annual Research Review: Childhood maltreatment, latent vulnerability and the shift to preventative psychiatry – the contribution of functional brain imaging

Eamon J. McCrory,^{1,2} Mattia I. Gerin,^{1,2} and Essi Viding¹

- Threat processing
 - i. Neural correlates
- Emotion Regulation
 - ii. Association with psychiatric disorder
- Reward Processing
 - iii. Association with maltreatment
- Executive Functioning

Annual Research Review: Childhood maltreatment, latent vulnerability and the shift to preventative psychiatry – the contribution of functional brain imaging

Eamon J. McCrory,^{1,2} Mattia I. Gerin,^{1,2} and Essi Viding¹

- Neurocognitive changes are observable even in the absence of psychiatric disorder and in some cases, predict future symptomatology. They are thought, in part, to reflect adaptations to early adverse environments.
- These changes are strikingly consistent with those seen in individuals presenting with psychiatric disorder suggesting such neurocognitive 'adaptations' embed latent vulnerability to future psychiatric disorder.

1. Altered threat processing

Children exposed to physical maltreatment have been shown to have altered processing of angry faces:

- able to more accurately identify angry facial expressions using sparse perceptual information than peers
- devote more attentional resources to the processing of angry faces
 - interpreted as increased hypervigilance to threat



Contents lists available at ScienceDirect

Cognition

journal homepage: www.elsevier.com/locate/COGNIT



Brief article

Development of perceptual expertise in emotion recognition


Seth D. Pollak^{a,*}, Michael Messner^a, Doris J. Kistler^b, Jeffrey F. Cohn^c

^aDepartment of Psychology and Waisman Center, University of Wisconsin at Madison, 1500 Highland Avenue, Madison, WI 53711, USA

^bUniversity of Louisville, The Heuser Hearing Institute, 117 E. Kentucky Street, Louisville, KY 40203, USA

^cUniversity of Pittsburgh, 4327 Sennott Square, Pittsburgh, PA 15260, USA





1. Altered threat processing

Children exposed to physical maltreatment have been shown to have altered processing of angry faces:


- able to more accurately identify angry facial expressions using sparse perceptual information than peers
- devote more attentional resources to the processing of angry faces - interpreted as increased hyper-vigilance to threat
- In some contexts they show **avoidance** of threat cues – diverting attention away from threat cues that may be processed as aversive

Current Biology

Heightened neural reactivity to threat in child victims of family violence

Dec 2011

Eamon J. McCrory^{1,2,*},
Stéphane A. De Brito^{1,2,*},
Catherine L. Sebastian¹,
Andrea Mechelli³, Geoffrey Bird^{4,5},
Phillip A. Kelly^{1,2}, and Essi Viding¹



Current Biology

Heightened neural reactivity to threat in child victims of family violence

Dec 2011

Eamon J. McCrory^{1,2,*},
Stéphane A. De Brito^{1,2,*},
Catherine L. Sebastian¹,
Andrea Mechelli³, Geoffrey Bird^{4,5},
Phillip A. Kelly^{1,2}, and Essi Viding¹





Current Biology

Heightened neural reactivity to threat in child victims of family violence

Dec 2011


Eamon J. McCrory^{1,2,*},
Stéphane A. De Brito^{1,2,*},
Catherine L. Sebastian¹,
Andrea Mechelli³, Geoffrey Bird^{4,5},
Phillip A. Kelly^{1,2}, and Essi Viding¹





y = 2

Increased right **amygdala** reactivity and increased bilateral **anterior insula** reactivity to angry vs. calm faces in children exposed to family violence.



z = -2

z = 4


This may be a latent neural marker of latent vulnerability – the same neural signature is common in anxiety disordered populations (Etkin & Wager, 2007).

Exposure to family violence may ‘recalibrate’ responsiveness of the anterior insula and amygdala in processing potential threat.

But is this a conscious process? In other words, is this hypervigilance to threat under higher order regulatory influence?

Amygdala activation in maltreated children during pre-attentive emotional processing

Eamon J. McCrory, Stéphane A. De Brito, Philip A. Kelly, Geoffrey Bird, Catherine L. Sebastian, Andrea Mechelli, Sophie Samuel and Essi Viding



Congruent

Duration
(ms)

+

500

*

1100

Congruent

Duration
(ms)



500



17




68



1100

Angry > Neutral




Amygdala

Y = -1


Is increased amygdala reactivity an adaptive response to environmental threat?



Children



Amygdala




McCrory et al., (2011)

Soldiers




Anterior
insula




Wingen et al., (2011)

Prior amygdala reactivity to threat cues predicts anxiety and depression symptoms in a cohort of health adults (n=340) following future life stressors over a 1 – 4 year period




Adversity



Altered Threat
Processing

- Increasing likelihood of interpersonal conflict with peers




Social Environment

- Reducing cognitive capacity available for more normative developmental tasks and social learning



Psychiatric Vulnerability



Autobiographical Memory (ABM)

Autobiographical memory

- Autobiographical memory (ABM) is concerned with the recollection of personally experienced events and plays a central role in scaffolding our sense of self and our ability to remain oriented in the present
- A constructive, flexible ABM provides the 'data' that helps us simulate future events and negotiate them more effectively (*The Constructive Episodic Simulation Hypothesis*, Schacter & Addis, 2007).
- It is thought that episodic simulation has particular adaptive value because it allows us to simulate a variety of ways in which the future might unfold without having to engage in actual behaviour (cf., Ingvar, 1979; Schacter, 2012; Suddendorf & Corballis, 1997, 2007).
- If ABM is **OVERGENERAL** then memories are characterized by greater categorical recollection and a paucity of specific detail.

Autobiographical memory

- Over-general ABM is associated with *current* psychopathology:
 - Depression (Sumner et al., 2010)
 - PTSD (Ono et al., 2016)
 - Schizophrenia (e.g. McDougall et al., 2015)
- Overgeneral ABM *predicts* symptoms of **Depression** in at-risk adolescents (e.g. Rawal & Rice, 2015). It also *predicts* symptoms of **PTSD** in assault survivors 6 months later even taking into account assault severity and baseline symptoms (Kleim & Ehlers, 2008).
- These findings are consistent with the notion that OGM is implicated in the **pathogenesis** of psychiatric disorder.
- Maltreatment is reliably associated with OGM at the behavioural level (Valentino et al., 2009; see Hitchcock et al., 2014 for a review).

Autobiographical memory: a candidate latent vulnerability mechanism for psychiatric disorder following childhood maltreatment

Eamon J. McCrory, Vanessa B. Puetz, Eleanor A. Maguire, Andrea Mechelli, Amy Palmer, Mattia I. Gerin, Philip A. Kelly, Iakovina Koutoufa and Essi Viding

BJPsych The British Journal of Psychiatry


Sept 2017

- The Autobiographical Memory Test (AMT), a standard measure of OGM, was administered to all participants in a session 1-3 weeks before scanning. Participants generated specific memories in response to 10 positive (e.g. achieve, caring) and 10 negative cue words (e.g. mistake, lonely). OGMs were defined as *'memories that did not contain at least one specific detail that identifies an event as a distinct episode'*.

| | Maltreatment Group (N=34) | Non-Maltreatment Group (N=33) | <i>p</i> |
|-----------------------|---------------------------|-------------------------------|----------|
| AMT Total | 17.4% | 10.0% | .017 |
| AMT Positive Memories | 15.8% | 9.6% | .066 |
| AMT Negative Memories | 19.0% | 10.4% | .018 |


Autobiographical memory

- The maltreated vs. non-maltreated group showed DECREASED activation to **positive** memories in the hippocampus
- This was interpreted as possibly reflecting reduced memory specificity of positive memories in maltreated children
- SDQ Total Score correlated positively with hippocampal ($r=0.47$, $p=0.007$) activation during negative memory recall in the Maltreated group (*greater specification of negative memories?*)




Autobiographical memory

- The maltreated vs. non-maltreated group showed INCREASED activation to **negative** memories in the amygdala and increased connectivity between the amygdala and the mACC indicating heightened salience processing
- This was interpreted as possibly reflecting a privileging of negative memories in the maltreated children.
- SDQ Total Score correlated positively with amygdala ($r=0.36$, $p=0.049$) activation during negative memory recall in the maltreated group




Autobiographical memory

- In a study of depressed patients, remitted patients and individuals at high familial risk of depression, Young and colleagues (2016) reported increased amygdala activation, and functional connectivity with regions implicated in salience processing (including the dorsal anterior cingulate cortex) during negative ABM recall in all three groups relative to healthy controls.
- This suggests that heightened amygdala activation and functional connectivity with the salience network during negative ABM recall may therefore represent a trait-like marker of depression.





- Poorer social problem solving
-> greater peer problems
- Development of negative self schema




- Greater ruminative processing
- Low mood

Adversity



Over-general
Autobiographical
Memory

- Poorer social problem solving – greater peer problems




Social Environment

- Increased negative ruminative style
- Poorer ability to conceptualize the future self




Psychiatric Vulnerability




Summary

- Relatively recent fMRI research has demonstrated that childhood maltreatment is associated with altered functioning in a range of neurocognitive systems including: threat processing and autobiographical memory processing. Reward processing is also implicated.
- Such changes are observable even in the absence of psychiatric disorder and in some cases, predict future symptomatology. They are thought, in part, to reflect adaptations to early adverse environments.
- These changes are strikingly consistent with those seen in individuals presenting with psychiatric disorder suggesting such neurocognitive 'adaptations' embed latent vulnerability to future psychiatric disorder.
- These findings establish a compelling case to develop a more precise mechanistic understanding of the pathogenesis of psychiatric disorder following maltreatment and the need to invigorate efforts to build a preventative clinical approach.
- Need to move on from meta-level clinical constructs such as 'trauma-focussed' and 'attachment-focussed'. Both are important, but primarily signpost the direction of travel; more precision needed as to the targets and mechanisms of change.



Infancy.....Childhood.....Adolescence.....Adulthood



Outcome

Unhealthy


Healthy

Clinical Threshold

Maltreatment



Infancy.....Childhood.....Adolescence.....Adulthood



Thank-you!



Developmental
Risk and Resilience Unit



Anna Freud
National Centre for
Children and Families

NSPCC

E·S·R·C
ECONOMIC
& SOCIAL
RESEARCH
COUNCIL